



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

LXXXI. *Some Considerations on a late Treatise intituled, A new Set of Logarithmic Solar Tables, &c. intended for a more commodious Method of finding the Latitude at Sea, by Two Observations of the Sun; by H. Pemberton, M. D. R. S. Lond. et R. A. Berol. S.*

Read Nov. 20,
1760.

AS it happens not unfrequently at sea for the unseasonable intervention of clouds to prevent the ordinary method of determining the ship's latitude by the sun's meridian altitude, even when it is of primary consequence, that the true latitude should be known; a problem for remedying this disappointment is stated in many treatises of navigation, for finding the latitude of a place by any two altitudes of the sun, with the interval of time between them.

A problem similar to this is proposed, and solved instrumentally upon a globe, by a very early writer, Petrus Nonius (*a*), namely, to find the latitude by two altitudes of the sun, and the angle made by the azimuth circles passing through the sun, when the altitudes are taken. And since more commodious and accurate instruments for measuring time have been invented, than were known to this author, the other problem has been proposed for the same pur-

(*a*) In Libr. De Observ. Regul. et Instrum. Geometr. lib. ii. c. 14.





pose, of which a construction upon the principles of the stereographic projection of the sphere is exhibited by Mr. Collins, in his Mariner's Plain Scale new planed (*b*). And as the direct method of solving both these problems by numbers requires a diversity of trigonometrical operations, a set of tables has lately been published for a more compendious way of computation in the problem, where the interval of time is given, whereby the ship's true latitude may be very expeditiously derived from the ship's dead reckoning, provided the observations are made within certain limits of time.

But however worthy of notice this method may be, new tables for the purpose are altogether unnecessary. It consists of two parts: the first computes, from the latitude exhibited by the dead reckoning of the ship, the distance from noon of the middle time between the observations, and thence the time of either: the second operation computes, from one of these observations, what should be the sun's meridian altitude, had the ship's reckoning given the true latitude; but if the latitude assumed from that reckoning is erroneous, the altitude thus computed will not be conformable to it; however, if the times for the observations are properly chosen, it will much better agree to the true latitude, and thereby the assumed latitude may be more or less corrected.

But both these operations are an immediate consequence from the proposition in spherical trigonometry, usually delivered under the name of the fourth axiom, which is this; That the square of the radius is to the rectangle under the sines of the sides con-

(*b*) Book iii. p. 35.

taining any angle, as the verfed line of that angle to the difference between the verfed lines of the third fide, and of the difference between the fides containing the angle.



Hence if A be the zenith of any place; B the elevated pole; C, D two places of the fun in the parallel of declination DF; by this proposition in the triangle ABC,

Rad.⁹ : fin. AB × fin. BC (cof. lat. × cof. decl. ⊙)
 ∴ verf. f. ABC : verf. f. AC — verf. f. AF;

And, in like manner, in the triangle ABD,
 Rad.⁹ : cof. lat. × cof. decl. ⊙ ∴ verf. f. ABD
 : verf. f. AD — verf. f. AF;

Consequently,

Rad.⁹ : cof. lat. × cof. decl. ⊙ : : verf. f. ABD — verf. f. ABC
 : verf. f. AD — verf. f. AC.

But $\frac{1}{2}$ rad. × $\overline{\text{verf. f. ABD} - \text{verf. f. ABC}}$ is =
 fin. $\frac{ABD + ABC}{2}$ × fin. $\frac{ABD - ABC}{2}$;

Therefore,

Rad.⁹ : cof. lat. × cof. decl. ⊙ : : fin. $\frac{ABD + ABC}{2}$
 × fin. $\frac{ABD - ABC}{2}$: $\frac{1}{2}$ rad. × $\overline{\text{verf. f. AD} - \text{verf. f. AC}}$.

And this is the first operation in the treatise, these remarks concern. For in the rectangle, the fin. $\frac{ABD + ABC}{2}$
 × the fin. $\frac{ABD - ABC}{2}$, the first side, when the observations are both made on the same side of noon, and the second side of this rectangle, when one observation is before noon, and the other after, is the sine of the distance of the middle time between the observations from noon, and the other side is the sine of half the distance between the observations. And in the new tables one column exhibits the arithmetical complement of the logarithmic sine of half the elapsed time, and the next, the logarithm of twice the sine of the middle time.

But moreover, $\frac{1}{2}$ rad. × $\overline{\text{verf. f. AD} - \text{verf. f. AC}}$
 being equal to fin. $\frac{AD + AC}{2}$ × fin. $\frac{AD - AC}{2}$, the operation may be expressed more commodiously thus,
 Rad.

$$\text{Rad.}^s : \cos. \text{ lat. } \times \cos. \text{ decl. } \odot :: \sin. \frac{\text{ABD} + \text{ABC}}{2} \\ \times \sin. \frac{\text{ABD} - \text{ABC}}{2} : \sin. \frac{\text{AD} + \text{AC}}{2} \times \sin. \frac{\text{AD} - \text{AC}}{2} ;$$

now requiring the common logarithmic fines only.

Again, the square of the radius being, as above, to $\cos. \text{ lat. } \times \cos. \text{ decl. } \odot$, as the $\text{vers. f. } \text{ABC}$ to $\text{vers. f. } \text{AC}$ — $\text{vers. f. } \text{AF}$, or as $\text{vers. f. } \text{ABD}$ to $\text{vers. f. } \text{AD}$ — $\text{vers. f. } \text{AF}$, these analogies express the second operation, which, in this treatise, is unnecessarily confined to the lesser angle ABC . The column in the table intituled *Rising* consists of logarithmic versed fines, which may be applied to either of the angles ABC or ABD promiscuously; for here AF being equal to the excess of BC or BD above the arch BA , assumed for the complement of the latitude, the arches AC and AD , in the preceding analogies, will be the complements of the altitudes observed, if the latitude were truly assumed, otherwise not; but the difference of their versed fines, will however be equal to the difference of the versed fines of the complements of the true latitudes; for this is supposed in the first operation. Therefore, if one of the fourth terms of these analogies be deducted from the versed fine of the complement of the greater altitude, or the other from the versed fine of the complement of the lesser, the remainder will be the same, and exhibit a versed fine for the complement of the sun's altitude different from AF , when BA is assumed different from the true latitude, and nearer to the truth, if the times for the observations are properly chosen.

But farther, the two preceding analogies may be reduced to these;

Rad.

$$\text{Rad.}^q : \frac{\text{cof. lat.} \times \text{cof. decl. } \odot}{: \frac{1}{2} \text{ rad.} \times \text{verf. f. AC} - \text{verf. f. AF}}, \quad \overline{\text{fin. } \frac{1}{2} \text{ ABC}}^q$$

And

$$\text{Rad.}^q : \frac{\text{cof. lat.} \times \text{cof. decl. } \odot}{: \frac{1}{2} \text{ rad.} \times \text{verf. f. AC} - \text{verf. f. AF}}, \quad \overline{\text{fin. } \frac{1}{2} \text{ ABD}}^q$$

Also,

$$\text{Verf. f. AC} - \text{verf. f. AF} = \text{cof. AF} - \text{cof. AC}, \text{ and} \\ \text{verf. f. AD} - \text{verf. f. AF} = \text{cof. AF} - \text{cof. AD}.$$

And thus the common tables of fines will supply the use of this column of the new tables, as well as the preceding.

The first example of this treatise, wrought by the common tables.

Here the declination of the sun is stated at $11^{\circ} 17'$ N. the altitude first taken, being the lesser, $46^{\circ} 55'$, the greater altitude $54^{\circ} 7'$, and the difference of time between the observations $1^h 25^m$, or 85^m ; which, divided by 4, gives $21^{\circ} 15'$ for the arch of the equinoctial corresponding.

Here the observations are both before noon; therefore, in fig. 1. the latitude being assumed $46^{\circ} 50'$,

For the first operation.

	AC =	35	53	0	
	AD =	43	5	0	
	Their sum =	78	58	0	
	Their difference =	7	12	0	
	Sine of $\frac{1}{2}$ sum (39 29 0)				9.80336
	Sine of $\frac{1}{2}$ difference (3 36 0)				8.79789
Arith. com. of the sine of $\frac{1}{2}$ CBD (= 10 37 30)					0.73429
Arith. com. of the sine of BF, or the cof. of the decl. of the sun, - - - - -					0.00848
				Sum	19.34402
	Cof. of the lat. assumed				9.83513
Diff. = fin. ABD + ABC					9.50889
				2	

[916]

For the second operation.

$$\frac{ABD + ABC}{2} = 18^{\circ} 49' 50''$$

$$\frac{ABD - ABC}{2} = 10^{\circ} 37' 30''$$

$$ABC = 8^{\circ} 12' 20''$$

$$\text{Sine of } \frac{1}{2} ABC (4^{\circ} 6' 10'') \left\{ \begin{array}{l} 8.85458 \\ 8.85458 \end{array} \right.$$

$$\text{Cof. of decl. } \odot \quad 9.99152$$

$$\text{Cof. of the lat. assumed} \quad 9.83513$$

$$\text{Sum} - 30.00000 = \log. \text{ of the } \frac{\text{verf. f. AC} - \text{verf. f. AF}}{2} \quad 7.53581$$

$$\begin{array}{r} \text{Verf. f. AC} - \text{verf. f. AF} \quad 0.006868 \\ \text{Natural cof. of AC, the complement of the greater} \\ \text{altitude, - - - - -} \quad \left\{ \begin{array}{l} 0.810212 \\ 0.817080 \end{array} \right. \\ \text{Natural cof. of } 35^{\circ} 12' 23'' \end{array}$$

$$\text{Or, } \frac{ABD + ABC}{2} + \frac{ABD - ABC}{2} \text{ being} = ABD,$$

$$\text{Sine of } \frac{1}{2} ABD (14^{\circ} 43' 40'') \left\{ \begin{array}{l} 9.40522 \\ 9.40522 \end{array} \right.$$

$$\text{Cof. of the decl. of the sun} \quad 9.99152$$

$$\text{Cof. of the latitude assumed} \quad 9.83513$$

$$\text{Sum} - 30.00000 = \log. \text{ of the } \frac{\text{verf. f. AD} - \text{verf. f. AF}}{2} \quad 8.63709$$

$$\begin{array}{r} \text{Verf. f. AD} - \text{verf. f. AF} \quad 0.086720 \\ \text{Natural cof. of AD, the complement of the lesser} \\ \text{altitude, - - - - -} \quad \left\{ \begin{array}{l} 0.730361 \\ 0.817081 \end{array} \right. \\ \text{Natural cof. of } 35^{\circ} 12' 23'', \text{ as before} \end{array}$$

Thus the meridian distance of the sun from the zenith is computed at $35^{\circ} 12' 23''$, and this added to the sun's declination $11^{\circ} 17' \text{ N.}$ gives $46^{\circ} 29' 23''$ for the latitude, different from that assumed, and nearer to the true latitude; which, by a second operation

operation with this latitude now found, will approach still nearer to the truth, thus:

The sum of the four first logarithms in the first of } 19.34402
the two former operations, - - - - - }

Cofine of $46^{\circ} 29' 23''$	9.83789
Sine of $\frac{ABD + ABC}{2} = 18^{\circ} 42' 25''$	9.50613
$\frac{ABD - ABC}{2} = 10^{\circ} 37' 30''$	
$ABC = 8^{\circ} 4' 55''$	
$\frac{1}{2} ABC = 4^{\circ} 2' 27\frac{1}{2}''$	$\left\{ \begin{array}{l} 8.84800 \\ 8.84800 \\ 9.99152 \\ 9.83789 \end{array} \right.$
Log. $\frac{\text{verf. f. AC} - \text{verf. f. AF}}{2}$	7.52541
Verf. f. AC — verf. f. AF	0.006704
Natural cof. of AC, as before,	0.810212
Natural cof. of $35^{\circ} 13' 21''$	0.816916

The declination of the sun added to this, amounts to $46^{\circ} 30' 31''$, the true latitude, as directly computed, being $46^{\circ} 30' 19''$.

However, it must not be expected, that this method of computation will always converge thus expeditiously to the true latitude. Had these observations been made about 3 hours sooner in the day, and with the same interval between them, if the first altitude of the sun had been found $18^{\circ} 55'$, and the second $33^{\circ} 11'$, the latitude computed would come out wider from the truth, than that assumed, one exceeding, and the other falling short of the true latitude, which will lie between them, and is nearly the same, as above. But if the lesser altitude had

been $29^{\circ} 13'$, and the greater 40° ; in 9 degrees of north latitude the interval between the observations would have been nearly the same, as before. But if this latitude be sought by any latitude assumed near it, the latitude computed according to the method above, will fall more remote from the truth, than that assumed, and err the same way.

In general, the error in the latitude assumed will bear to the error in the latitude computed, nearly the ratio compounded of that of the rectangle under the radius and the cosine of the distance from noon of the middle point of time between the two observations to twice $\sin. \frac{1}{2} ABC \times \sin. \frac{1}{2} ABD$, and the ratio of $\text{rad.} \times \sin. AF$ to $\sin. BC \times \cos. AB$; inasmuch that the distance of the middle point of time between the two observations from noon is to be considered, as the limit, where these computations shall cease to converge, when the rectangle under the radius, and the cosine of this distance from noon shall be to twice $\sin. \frac{1}{2} ABC \times \sin. \frac{1}{2} ABD$, (or the cosine of this distance to the difference between the versed sine of this distance from the versed sine of half CBD) as $\sin. BF \times \cos. AB$ to $\text{rad.} \times \sin. AF$. And the errors in the assumed and computed latitudes fall on different sides of the true latitude, when both the observations are on the same side of noon, and the zenith lies between the meridian sun and the elevated pole; or when the observations are one before, and the other after noon, if the meridian sun passes between the zenith and the elevated pole: otherwise, they fall on the same side of the true latitude.

However,

However, the direct method of computation, by the assistance of the natural sines, will not be so much more operose than this compendium, as may at first sight be imagined. For the arch of a great circle being drawn through C and D, forming the triangle BCD, if the logarithmic cosine of BC or BD be added to the logarithmic tangent of half the angle CBD, and the logarithmic sine of BC or BD be added to the logarithmic sine of half this angle, the first sum is the logarithmic cotangent of the angle BCD, and the second the logarithmic sine of half CD, the base of the triangle. Then, in the triangle ACD, from the sides, now all given, is to be computed the angle ACD, the difference between which, and the angle above found BCD, is the angle BCA, when the zenith lies between the pole, and the great circle through C, D; but when the zenith lies beyond that circle, the angle BCA is either the sum of those angles, or the supplement of that sum to a circle. And, in the last place, if twice the logarithmic sine of half this angle, and the logarithmic sines of BC and AC are added together, the sum, after thrice the logarithm of the radius has been deducted, is the logarithm of half the excess of the natural cosine of BC ∞ AC above the natural cosine of AB, or the natural sine of the latitude, according to the trigonometrical axiom, which has been above referred to; for $\text{rad.} \times \frac{1}{2} \text{ vers. f. } BCA$ is equal to $\text{fin. } \frac{1}{2} BCA^{\text{a}} (c)$.

(c) Vide Neper. Mirif. Canon. Construct. Edinburg. 1619, p. 59.

The operation for the preceding example will be this :

<p>Tang. $\frac{1}{2}$ CBD — 9.273225 Cof. BC — 9.291504</p> <hr style="width: 100%;"/> <p>Cotang. BCD — 9.564729</p> <hr style="width: 100%;"/> <p style="text-align: center;">BCD = $87^{\circ} 53' 52''$</p>	<p>Sine $\frac{1}{2}$ CBD — 9.265714 Sine BC — 9.991524</p> <hr style="width: 100%;"/> <p>Sine $\frac{1}{2}$ CD — 9.257238</p> <hr style="width: 100%;"/> <p style="text-align: center;">$\frac{1}{2}$ CD = $10^{\circ} 25' 2\frac{1}{2}''$</p>
--	--

<p>AD = $43^{\circ} 5' 0''$ AC = $35^{\circ} 53' 0''$ CD = $20^{\circ} 50' 5''$ Sum = $99^{\circ} 48' 5''$ $\frac{1}{2}$ Sum = $49^{\circ} 54' 2\frac{1}{2}''$</p>	<p>0.232001 } ar.co. of the fines. 0.448950 } 9.883621 } fines. 9.074466 } Sum — 19.639038 $\frac{1}{2}$ Sum — 9.819519</p> <hr style="width: 100%;"/>
<p>$\frac{1}{2}$ Sum — AD = $6^{\circ} 49' 2\frac{1}{2}''$</p>	

This half sum is the logarithmic cofine of half ACD; therefore, half ACD is = $48^{\circ} 42' 11''$, and its difference from $\frac{1}{2}$ BCD will give $\frac{1}{2}$ ACB = $4^{\circ} 45' 15''$.

Sine of $\frac{1}{2}$ ACB	$\left\{ \begin{array}{l} 8.918444 \\ 8.918444 \end{array} \right.$
Sine of AC —	9.767999
Sine of BC —	9.991524
	<hr style="width: 100%;"/>
0.0039483 — log. —	7.596411

0.0078966

0.7333345 — natural cofine of BC — AC, that is, of $42^{\circ} 50'$.


0.7254379 — natural cofine of AB, or fine of the latitude.

Therefore, the latitude is $46^{\circ} 30' 19''$.

But farther, as in the cases, where the above approximation takes place, it is an advantage, that the assumed latitude should be taken as near the true one,

one, as can be, the dead reckoning of the ship need not be intirely relied on for that purpose; for the instrumental construction proposed by Mr. Collins (*d*) will very readily give the true latitude, as nearly as the instruments used can express: and his method may be described thus.

Two straight lines AB, AC being drawn in an angle corresponding with the distance in time between the observations, and in one of the lines, as AB, the lengths AD, AE being taken from any scale of tangents, one equal to the tangent of half the sum, and the other equal to the tangent of half the difference of the distance of the sun from the pole, and from the zenith of the first observation, and also the points F, G, taken in the other line in the same manner related to the other observation; then circles being described on the two diameters DE, FG, the distance of A from one of the intersections of these circles will be the tangent of half the complement of the latitude.



Moreover, as in the treatise, which has given occasion to this discourse, it is proposed sometimes to take into consideration the ship's motion during the time, between the observations, but imperfectly, regard being had to the change made by the ship in longitude only; but the change in latitude also may be taken into consideration in the foregoing con-

(*d*) In the place above cited.

struction, by making the angle BAC equal to the sum or difference of the interval of time, and the change in the ship's longitude, according as that change is made towards the west, or towards the east, and then drawing a line from A , not to the intersection of the circles; but so, that the portions of that line, terminated by each of the circles, may be the tangents of arches, whose difference is half the change in the latitude; which, if made from the pole, requires the portion of the line terminated at the circle, whose diameter is DE , to be shorter than the other; and the contrary is required, when the motion of the ship is towards the pole.



For determining the latitude by calculation, if the distance of the sun in the first observation from the zenith of the second could be found, this case would be reduced to the first, wherein the ship is considered, as stationary. And for this purpose, it has been proposed to make an additional observation, by an azimuth compass, of the angle, the ship's course makes with the azimuth of the sun, when the first altitude is taken; and, perhaps, the same angle may be found with sufficient exactness from the latitude in the first observation assumed, whether from the dead reckoning, or from the foregoing construction.

Why I have taken no notice of the calculations exhibited by the author of the piece here animadverted on, as proofs of his method, will readily appear to those, who shall cast their eye upon them.

SCHOLIUM.


The axiom in trigonometry, on which the calculations here discoursed of have been shewn immediately

diately to depend, was introduced by Regiomontanus, and is still retained, as the foundation of the present methods of computing logarithmically an angle from the three sides of a spherical triangle given, though they may be demonstrated more directly by the following lemma.



In the circle ABC, the chord AC being drawn, and the arch ABC bisected in B, if D be taken in the circle at pleasure, and the lines BE, DF drawn also at pleasure parallel to each other, one terminated by the circle in E, and the other by the chord AC in F; then AD, DC being drawn, $BE \times DF = AD \times DC$.

Draw DH perpendicular to AC, and also the diameter BG, joining EG. Then the angles FDH, EBG are equal, and the angles DHF, BEG right. Therefore BE is to BG as DH to DF, and $BE \times DF = BG \times DH = AD \times DC$.



Now, in the spherical triangle ABC , through B , and to the poles A, C , let the semicircles EBF, GBD be described, and the arch AC completed to a circle. Then the semicircles EBF, GBD will be both perpendi-

cular to the plane of this circle, cut it in their diameters EF, GD , and their common intersection BH be also perpendicular to it, and consequently perpendicular to GD , the diameter of the semicircle GBD , whereby GB, DB being joined, $GD \times GH$ is equal to GB^2 , and $GD \times DH$ equal to BD^2 . Again, AK being drawn parallel to GD , the arch ACK is double AC , GCD double CB , EAD the sum of all the three sides of the triangle, $EG = ED - DG$ ($2CB$), $GF = AF$ (AE) $+ CG$ (CD) $- AC = ED - 2AC$, and $FD = ED - EF$ ($2AB$).

Then EG, GF, FD, ED being joined; in the first place, AK is to DG as $AK \times GH$ to $DG \times GH$, also AK to DG as $AK \times DG$ to DG^2 , but $AK \times GH = EG \times GF$ by the preceding lemma, and $DG \times GH = GB^2$; whereby $AK \times DG$ is to DG^2 as $EG \times GF$ to GB^2 . Hence AK being twice the sine of the arch AC , DG twice the sine of the arch GC , equal to BC , EG twice the sine of half the arch EAG , GF twice

twice the fine of half the arch GF, and GB twice the fine of half the angle ACB to the radius of the circle GBD, which is half GD; $\text{fin. AC} \times \text{fin. BC}$ is to rad.^q as $\text{fin. } \frac{1}{2} \text{ED} - \text{BC} \times \text{fin. } \frac{1}{2} \text{ED} - \text{AC}$ to $\text{fin. } \frac{1}{2} \text{ACB}^q$. And this is Napeir's first method of finding an angle from the three sides given (*e*), as it is usually delivered.

Again, AK is to DG, or $\text{AK} \times \text{DG}$ to DG^q , as $\text{AK} \times \text{DH}$ ($\text{ED} \times \text{DF}$) to $\text{DG} \times \text{DH}$, or DB^q . Therefore, DB being twice the fine of half the angle BCD, or twice the cofine of half the angle ACB to the radius of the circle GBD, as $\text{fin. AC} \times \text{fin. CB}$ to rad.^q so is $\frac{1}{2} \text{ED} \times \text{fin. } \frac{1}{2} \text{ED} - \text{AB}$ to $\text{cof. } \frac{1}{2} \text{ACB}^q$. And this is Napeir's second method (*f*).

And lastly, the cofine of an arch or angle being to its fine, as radius to the tangent, $\text{fin. } \frac{1}{2} \text{ED} \times \text{fin. } \frac{1}{2} \text{ED} - \text{AB}$ will be to $\text{fin. } \frac{1}{2} \text{ED} - \text{BC} \times \text{fin. } \frac{1}{2} \text{ED} - \text{AC}$ as rad.^q to $\text{tang. } \frac{1}{2} \text{ACB}^q$, which is a third method added by Gellibrand (*g*).


Moreover,

In plain triangles, from the three sides given may an angle be found by a process similar to each of these, as follows:

(*e*) Mirific. Canon. Log. descrip. lib. ii. c. 6. § 8.

(*f*) Ibid. § 9.

(*g*) Trigon. Britan. p. 75.



In the plain triangle ABC , if on the centers A and C be described circles through B , cutting AC in E, F, G , and D , the perpendicular BH being let fall on AC , $EH \times HF$ is equal to $GH \times HD$. Therefore $HF : GH :: HD : HE$, and by composition $GF : GH :: ED : HE$, and $:: ED - GF : EG$; also $HF : GF :: HD : ED$, and $:: DF : ED - GF$.

Moreover ED is $= AB + AC + CB$, $EG = ED - 2CB$, $FD = ED - 2AB$, $ED - GF = EG + FD = 2ED - 2CB - 2AB = 2AC$. Hence GF is to GH as $2AC$ to EG , whence $GF \times EG$ is $= 2AC \times GH$; and DH is to ED as DF to $2AC$, that is, $2AC \times DH$ will be $= ED \times DF$.

Thus $2AC : DG (:: 2AC \times DG : DG^2) :: 2AC \times GH (EG \times GF) : DGH (GB^2)$ and $AC \times CB : \text{rad.}^2 :: \frac{1}{2}ED - CB \times \frac{1}{2}ED - AC : \sin. \frac{1}{2}ACB]^2$.


Likewise $2AC : DG (:: 2AC \times DG : DG^2) :: 2AC \times DH (ED \times DF) : GDH (DB^2)$; therefore $AC \times CB : \text{rad.}^2 :: \frac{1}{2}ED \times \frac{1}{2}ED - AB : \cos. \frac{1}{2}ACB]^2$.

And in the last place $\frac{1}{2}ED \times \frac{1}{2}ED - AB : \frac{1}{2}ED - CB \times \frac{1}{2}ED - AC :: \text{rad.}^2 : \tan. \frac{1}{2}ACB]^2$.

This last proposition was first delivered by Gellibrand, and the other two not long after by Oughtred, in their respective treatises of Trigonometry.

BUT it may farther here be noted, that from two sides of a plain triangle, as AC , CB , with the angle between them ACB , the third side AB may be found either by the first or second of the foregoing proportions; for by the first may be found $EG \times GF$, that is, $AF^2 - AG^2$, and by the second $ED \times DF$, or $AD^2 - AF^2$; whence by a table of squares AF ($= AB$) may be readily found, for AG is equal to $AC - BC$, and $AD = AC + CB$.

In like manner in the spherical triangle from the sides AC , CB , and the angle ACB , may the third side be expeditiously found with the aid of a table of natural sines, by the axiom referred to in the beginning of this discourse, as modelled for this purpose by Napeir, in the proposition, which has been already referred to (*b*), whereby the axiom is reduced to this analogy,



$$\text{Rad.}^2 : \sin. AC \times \sin. CB :: \overline{\sin. \frac{1}{2} ACB}^2 \\ : \text{rad.} \times \frac{\text{verf. f. } AB - \text{verf. f. } AC \propto CB}{2} \quad (i).$$

MAY

(*b*) Mirif. Canon. Constr. p. 59.

(*i*) An example of this method has been given above in the triangle ABC in p. 920. And in default of a table of natural sines, the base AB might have been found thus. The sum of twice the logarithmic sine of $\frac{1}{2} ACB$, added to the logarithmic sines of AC

and

MAY it here be farther added, that thus all the cases of oblique spherical triangles may be solved without dividing them into rectangular? For when two sides and the angle included, or when two angles and the side between them are given, the two other angles or sides may be found by the proportions of Napeir and Briggs, whereby the other two angles or sides are found together (*k*); the proportions, when two sides with the including angle are given, being these,

$$\text{Cof. } \frac{AC + CB}{2} : \text{cof. } \frac{AC \propto CB}{2} :: \text{cotang. } \frac{1}{2} ACB \\ : \text{tang. } \frac{CAB + CBA}{2}.$$

And

$$\text{Sin. } \frac{AC + CB}{2} : \text{fin. } \frac{AC \propto CB}{2} :: \text{cotang. } \frac{1}{2} ACB \\ : \text{tang. } \frac{CAB \propto CBA}{2}.$$

The second of these has been demonstrated with great conciseness by Dr. Halley, from the principles of the stereographic projection of the sphere (*l*); and the first is derived by him from the second, but not

and BC amounted to 37.596411. The logarithmic sine of $\frac{AC \propto BC}{2}$ (9.562468) deducted from half this sum (18.798211)

leaves 9.235737. This number sought among the logarithmic tangents exhibits for its correspondent sine 9.229400, which being deducted from the foresaid half sum (18.79211) leaves 9.568805, which is the logarithmic sine of $21^{\circ} 44' 50\frac{1}{2}''$, equal to half AB.

(*k*) Vide Neper. Mirif. Canon. Confr. pag. penult.

(*l*) See Jones's Synopsis, p. 281.

altogether

altogether with the same success, by his not observing, that if the base and one of the sides be completed to semicircles, the second of the two proportions in the supplemental triangle thence formed leads directly to the first in the original triangle.

And to conclude, if two sides, and the angle opposite to one of them, or two angles with the side opposite to one, were given; when the other opposite part is found from the proportion between the sines of parts opposite, the remaining angle, or side, may be found by either of the two proportions foregoing.

LXXXII. *An Account of the Plants Halefia and Gardenia: In a Letter from John Ellis, Esq; F. R. S. to Philip Carteret Webb, Esq; F. R. S.*

Dear Sir,

Read Nov. 20, 1760. **Y**OU must have observed, that as the spirit of planting has increased in this kingdom, the study of botany has become more fashionable; the works of the celebrated Linnæus, heretofore looked on as capricious and strange, are now in the hands of every man, who wishes to study the order of nature.

The great variety of plants, which you have introduced into your garden from North America, as well as from many other parts of the world, must give you double pleasure, when you view them ranged in proper order, and judiciously named.

The